

# ECON 6170 Section 6

TA: Patrick Ferguson

October 11, 2024


## The Relative Topology

The Correspondence notes use the concepts of relative closedness and relative openness. Let  $S \subseteq \mathbb{R}^d$ .

**Definition 1.** A set  $U \subseteq S$  is open relative to  $S$  if every point  $x \in U$  is contained in a relatively open ball  $B_\varepsilon(x) \cap S \subseteq U$ .

**Example 1.**

- (i) Consider  $\mathbb{R}_+ \subseteq \mathbb{R}$ .  $[0, 1)$  is open relative to  $\mathbb{R}_+$ . For if  $x \in [0, 1)$  then  $(x - \varepsilon, x + \varepsilon) \cap \mathbb{R}_+ \subseteq [0, 1)$  for sufficiently small  $\varepsilon$ . In particular  $(-\varepsilon, \varepsilon) \cap \mathbb{R}_+ = [0, \varepsilon) \subseteq [0, 1)$ .
- (ii) Consider  $\mathbb{N} \subseteq \mathbb{R}$ .  $\{1\}$  is open relative to  $\mathbb{N}$ . For  $(1 - \varepsilon, 1 + \varepsilon) \cap \mathbb{N} = \{1\} \subseteq \{1\}$  for  $\varepsilon < 1$ .
- (iii) Consider  $X := \mathbb{R} \times \{0\} \subseteq \mathbb{R}^2$ .  $U := (a, b) \times \{0\}$  is open relative to  $X$ . For if  $x \in U$ , then  $B_\varepsilon(x) \cap X = (x - \varepsilon, x + \varepsilon) \times \{0\} \subseteq U$  for sufficiently small  $\varepsilon$ .



**Remark 1.** If it is clear we are using the relative topology of  $S$ ,  $B_\varepsilon(x)$  is often used in place of  $B_\varepsilon \cap S$ . For example in (i),  $B_\varepsilon(0)$  would be  $[0, \varepsilon)$ . This is the case in the Correspondence notes.

**Definition 2.** A set  $V \subseteq S$  is closed relative to  $S$  if for any sequence  $(x_n)$  in  $V$  that converges to  $x \in S$  has  $x \in V$ .

**Example 2.** Consider  $\mathbb{R}_{++} \subseteq \mathbb{R}$ .  $(0, 1]$  is closed relative to  $\mathbb{R}_{++}$ , because  $0 \notin \mathbb{R}_{++}$ .

**Remark 2.** If it is clear we are using the relative topology of  $S$ , if  $(x_n)$  is a sequence in  $S$  such that  $x_n \rightarrow x \notin S$ , we sometimes say that  $(x_n)$  diverges. That is, convergence means convergence to a point in  $S$ . In the previous example, the sequence  $(\frac{1}{n})$  would be said to diverge.

**Remark 3.** The following results about open and closed sets generalise to relatively open and relatively closed sets:

- $A$  is relatively open in  $S$  iff  $S \setminus A$  is relatively closed in  $S$ .
- The union of relatively open sets is also relatively open.
- The intersection of relatively closed sets is also relatively closed.
- The finite intersection of relatively open sets is also relatively open.
- The finite union of relatively closed sets is also relatively closed.

**Remark 4.** Some results don't generalise to the relative topology:

- Heine-Borel may not hold. That is, some sets that are bounded and relatively closed are not compact (in the sense of sequential or covering compactness). For example,  $(0, 1]$  is relatively closed in  $\mathbb{R}_{++}$  but  $\frac{1}{n}$  has no subsequence that converges in  $(0, 1]$ .
- Cauchy sequences may not converge. For example,  $(\frac{1}{n})$  is Cauchy but doesn't converge in the relative topology of  $\mathbb{R}_{++}$ .

**Remark 5.** Compactness is not relative: a set is (sequentially or covering) compact iff it is compact in every relative topology.

## Correspondences

**Definition 3.** We define the power set of  $X$  to be the set of all subsets of  $X$ , and denote it by  $2^X$ .

**Definition 4.** A correspondence  $F : X \rightrightarrows Y$ , from a nonempty set  $X$  to another nonempty set  $Y$ , is a function from  $X$  to  $2^Y$ .

- (i) The domain of  $F$  is  $X$ .
- (ii) The codomain of  $F$  is  $Y$ .
- (iii) The range of  $F$  is  $\bigcup \{F(x) \in 2^Y \mid x \in X\}$ .

In our course,  $X$  and  $Y$  will always be subsets of Euclidean space.

**Definition 5.** A correspondence is  $p$ -valued if the sets it maps to all have property  $p$ .

**Remark 6.** We can think of functions as a singleton-valued correspondences, in the sense that every function with values  $f(x)$  uniquely specifies a correspondence with values  $\{f(x)\}$ , and *vice versa*.

**Section Exercise 1.** Determine whether the following correspondences are closed-valued, compact-valued, singleton-valued, and/or convex-valued:

- (i) A budget correspondence,  $B : \mathbb{R}_{++}^{n+1} \rightrightarrows \mathbb{R}_+^n$ , given by  $B(p, w) := \{x \in \mathbb{R}_+^n \mid p \cdot x \leq w\}$ .  
 $p \cdot 0 \leq w$ , so  $0 \in B$ . For fixed  $p, w$  the set  $B(p, w)$  is the intersection of the closed half-space  $p \cdot x \leq w$ , with the nonnegative orthant  $\mathbb{R}_+^n$ . It is therefore the intersection of closed sets, and so closed itself. Given  $p \gg 0$  and  $x \geq 0$ , it is also bounded, and so compact. If  $p \cdot x \leq w$  and

$p \cdot y \leq w$ , then  $p \cdot (\alpha x + (1 - \alpha)y) \leq \alpha w + (1 - \alpha)w = w$ , so it is convex. Therefore,  $B$  is a nonempty-valued, compact-valued and convex-valued correspondence.

(a) What happens if we let some of the prices be 0?

$B$  is no longer bounded-valued, and thus no longer compact-valued.

(ii) A Walrasian demand correspondence,  $X : \mathbb{R}_{++}^{n+1} \rightrightarrows \mathbb{R}_+^n$ , given by  $X(p, w) := \arg \max\{u(x) \mid x \in B(p, w)\}$

(a) ... without restrictions on  $u$ .

$X(p, w) \subseteq B(p, w)$ , so  $X$  is bounded-valued. But it may be neither closed nor convex. Take, for example, the utility function  $u : \mathbb{R}_+^2 \rightarrow \mathbb{R}$  given by

$$u(x_1, x_2) = \mathbf{1}_{\{x_1 \neq x_2\}}$$

Then  $(0, \frac{1}{n})$  is a sequence that is (eventually) in  $X(p, w)$ , but the limit,  $(0, 0) \notin X(p, w)$ , so  $X$  is not closed-valued. Moreover,  $(\varepsilon, 0)$  and  $(0, \varepsilon)$  are in  $X(p, w)$  for sufficiently small  $\varepsilon > 0$ , but  $\frac{1}{2}(\varepsilon, 0) + \frac{1}{2}(0, \varepsilon) = (0, 0) \notin X(p, w)$ , so it is not convex-valued either.

(b) ... where  $u$  is continuous.

Continuity of  $u$  and compactness of  $B(p, w)$  imply that  $u$  attains a maximum on  $B(p, w)$ , so  $X$  is nonempty-valued. Let  $(x_i)$  be an arbitrary sequence in  $X(p, w)$ . Then  $u(x_i) = u^*$  for all  $i \in \mathbb{N}$ , so by continuity of  $u$ ,  $u(\lim x_i) = \lim u(x_i) = u^*$ . Therefore,  $X$  is closed-valued, and because it is bounded-valued, compact-valued.

(c) ... where  $u$  is continuous and quasiconcave.

Continuity implies that  $u$  attains a maximum on  $B(p, w)$ . Quasiconcavity implies  $X(p, w) = \{x \in B(p, w) \mid u(x) \geq \max_{y \in B(p, w)} u(y)\}$  is convex, so  $X$  is convex-valued.

(d) ... where  $u$  is continuous and *strictly* quasiconcave.

$X$  is singleton-valued. For if  $x \neq x'$  and  $u(x) = u(x') \geq u(y)$  for all  $y \in B(p, w)$ , then  $u(\frac{1}{2}x + \frac{1}{2}x') > u(x)$  by strict quasiconcavity, a contradiction.

(iii) An upper contour correspondence,  $R : \mathbb{R}_+^n \rightrightarrows \mathbb{R}_+^n$ , given by

$$R(x) := \{y \in \mathbb{R}_+^n \mid u(y) \geq u(x)\}$$

where  $u$  is continuous?

$R$  is nonempty-valued, for  $x \in R(x)$ .  $R$  is closed-valued, for if  $(y_i)_{i=1}^\infty$  is a convergent sequence in  $R(x)$ , then  $u(y_i) \geq u(x)$  for all  $i$ . By continuity of  $u$ , this implies that  $u(\lim_i y_i) \geq u(x)$ , so  $\lim_i y_i \in R(x)$ .

**Section Exercise 2.** Consider the following static game

	$L$	$R$
$U$	1, 1	0, 0
$D$	0, 1	1, 1

where player 1 chooses the row and player 2 chooses the column. The ordered-pair entries are the payoffs,  $(v_1, v_2)$ , to player 1 and player 2, respectively. Define the best response correspondence of

player 1  $BR_1 : \{L, R\} \rightrightarrows \{U, D\}$  by

$$x \in BR_1(y) \iff v_1(x, y) \geq v_1(x', y) \text{ for } x' \in \{U, D\}$$


Define the best response correspondence of player 2 similarly. Which player has a singleton-valued best response correspondence?

Player 1.  $BR_1(L) = \{U\}$  and  $BR_1(R) = \{D\}$ . In contrast,  $BR_2(D) = \{L, R\}$ .


**Definition 6.** The graph of a function  $f : X \rightarrow Y$  is the set of points  $\{(x, y) \in X \times Y \mid y = f(x)\}$ .

**Definition 7.** The graph of a correspondence  $F : X \rightrightarrows Y$  is the set of points  $\{(x, y) \in X \times Y \mid y \in F(x)\}$ .


**Remark 7.** As with functions, graphs of correspondences  $F : X \subseteq \mathbb{R} \rightrightarrows \mathbb{R}$  are easy to visualize. The difference is that for correspondences, a vertical line can intersect the graph more than once.




Singleton-valued




Compact- and convex-valued



Not convex-valued



Not convex-valued



Not closed-valued